

**QUICK INVENTORY OF DEPRESSIVE SYMPTOMATOLOGY (CLINICIAN-RATED)  
(QIDS-C)**

NAME: \_\_\_\_\_

TODAY'S DATE: \_\_\_\_\_

Please circle one response to each item that best describes the patient for the last seven days.

1. Sleep Onset Insomnia:

- 0 Never takes longer than 30 minutes to fall asleep.
- 1 Takes at least 30 minutes to fall asleep, less than half the time.
- 2 Takes at least 30 minutes to fall asleep, more than half the time.
- 3 Takes more than 60 minutes to fall asleep, more than half the time.

2. Mid-Nocturnal Insomnia:

- 0 Does not wake up at night.
- 1 Restless, light sleep with few awakenings.
- 2 Wakes up at least once a night, but goes back to sleep easily.
- 3 Awakens more than once a night and stays awake for 20 minutes or more, more than half the time.

3. Early Morning Insomnia:

- 0 Less than half the time, awakens no more than 30 minutes before necessary.
- 1 More than half the time, awakens more than 30 minutes before need be.
- 2 Awakens at least one hour before need be, more than half the time.
- 3 Awakens at least two hours before need be, more than half the time.

4. Hypersomnia:

- 0 Sleeps no longer than 7-8 hours/night, without naps.
- 1 Sleeps no longer than 10 hours in a 24 hour period (include naps).
- 2 Sleeps no longer than 12 hours in a 24 hour period (include naps).
- 3 Sleeps longer than 12 hours in a 24 hour period (include naps).

5. Mood (Sad):

- 0 Does not feel sad.
- 1 Feels sad less than half the time.
- 2 Feels sad more than half the time.
- 3 Feels intensely sad virtually all the time.

6. Appetite (Decreased):

- 0 No change from usual appetite.
- 1 Eats somewhat less often and/or lesser amounts than usual.
- 2 Eats much less than usual and only with personal effort.
- 3 Eats rarely within a 24-hour period, and only with extreme personal effort or with persuasion by others.

7. Appetite (Increased):

- 0 No change from usual appetite.
- 1 More frequently feels a need to eat than usual.
- 2 Regularly eats more often and/or greater amounts than usual.
- 3 Feels driven to overeat at and between meals.

8. Weight (Decrease) Within The Last Two Weeks:

- 0 Has experienced no weight change.
- 1 Feels as if some slight weight loss occurred.
- 2 Has lost 2 pounds or more.
- 3 Has lost 5 pounds or more.

9. Weight (Increase) Within the Last Two Weeks:

- 0 Has experienced no weight change.
- 1 Feels as if some slight weight gain has occurred.
- 2 Has gained 2 pounds or more.
- 3 Has gained 5 pounds or more.

**Enter the highest score on any 1 of the 4  
appetite/weight change items (6–9 above) \_\_\_\_\_**

**Enter the highest score on any  
1 of the 4 sleep items (1–4 above) \_\_\_\_\_**

10. Concentration/Decision Making:

- 0 No change in usual capacity to concentrate and decide.
- 1 Occasionally feels indecisive or notes that attention often wanders.
- 2 Most of the time struggles to focus attention or make decisions.
- 3 Cannot concentrate well enough to read or cannot make even minor decisions.

11. Outlook (Self):

- 0 Sees self as equally worthwhile and deserving as others.
- 1 Is more self-blaming than usual.
- 2 Largely believes that he/she causes problems for others.
- 3 Ruminates over major and minor defects in self.

12. Suicidal Ideation:

- 0 Does not think of suicide or death.
- 1 Feels life is empty or is not worth living.
- 2 Thinks of suicide/death several times a week for several minutes.
- 3 Thinks of suicide/death several times a day in depth, or has made specific plans, or attempted suicide.

13. Involvement:

- 0 No change from usual level of interest in other people and activities.
- 1 Notices a reduction in former interests/activities.
- 2 Finds only one or two former interests remain.
- 3 Has virtually no interest in formerly pursued activities.

14. Energy/Fatiguability:

- 0 No change in usual level of energy.
- 1 Tires more easily than usual.
- 2 Makes significant personal effort to initiate or maintain usual daily activities.
- 3 Unable to carry out most of usual daily activities due to lack of energy.

15. Psychomotor Slowing:

- 0 Normal speed of thinking, gesturing, and speaking.
- 1 Patient notes slowed thinking, and voice modulation is reduced.
- 2 Takes several seconds to respond to most questions; reports slowed thinking.
- 3 Is largely unresponsive to most questions without strong encouragement.

16. Psychomotor Agitation:

- 0 No increased speed or disorganization in thinking or gesturing.
- 1 Fidgets, wrings hands and shifts positions often.
- 2 Describes impulse to move about and displays motor restlessness.
- 3 Unable to stay seated. Paces about with or without permission.

**Enter the highest score on either of the 2 psychomotor items (15 or 16 above) \_\_\_\_**

Total Score: \_\_\_\_ (Range 0—27)